


Carbene-Catalyzed Indole 3-Methyl C(sp³)–H Bond FunctionalizationJian Cheng,[†] Jun Sun,[†] Jiekuan Yan,[†] Song Yang,^{*,†} Pengcheng Zheng,[†] Zhichao Jin,[†] and Yonggui Robin Chi^{*,†,‡}[†]Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering, Key Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, Guizhou University, Huaxi District, Guiyang 550025, China[‡]Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

Supporting Information

ABSTRACT: The metal-free catalytic functionalization of aromatic sp²-carbons and benzylic sp³-carbons remains challenging. Here we report a carbene-catalyzed functionalization of the 3-methyl sp³-carbon attached to 2-formyl-indoles. The reaction proceeds through an NHC-bound *o*-quinodimethane as the key intermediate generated from 2-formyl-3-methylindoles under oxidative conditions. Reactive ketones are found to be effective substrates to produce substituted hydropyrano[3,4-*b*]indoles in good to excellent yields.



INTRODUCTION

The indole rings are common fragments in natural products, agrichemicals, and pharmaceuticals.¹ Among the various indole-containing structures, hydropyrano[3,4-*b*]indole represents an important subtype core scaffold frequently found in bioactive molecules (Scheme 1a).² For example, Etodolac and its derivatives have been widely used as anti-inflammatory drugs,^{3a} potent analgesic agents,^{3b} and polymerase inhibitors.^{3c} Analogues and derivatives of hydropyrano[3,4-*b*]indoles also exhibit important bioactive properties and are frequently evaluated for biomedical applications.^{3d,e} Thus, synthetic methods that can effectively functionalize the sp² aromatic carbons and the sp³-carbons attached to the indole framework have received considerable attention. Progress in this area mainly comes from transition-metal-catalyzed C–H bond activations.⁴ Metal-free catalytic approaches are much less explored.⁵

Our laboratories are interested in developing metal-free N-heterocyclic carbene (abbreviated as NHC or carbene) organic catalysis for unique activation and reaction modes.⁶ To date, it has been established that multiple carbons (e.g., α -, β -, γ -, and δ -carbons) of (unsaturated) aldehydes can be efficiently activated via carbene organic catalysis (Scheme 1b).⁷ In contrast, functionalization of the sp²- and sp³-carbons of aromatic aldehydes remains difficult. We disclosed carbene-catalyzed functionalization of the 2-methyl sp³-carbon attached to heterocyclic aromatic aldehydes.⁸ Glorius and Rovis developed the functionalization of the sp³-carbon on 2-(bromomethyl)-benzaldehyde through NHC organocatalysis.⁹ Here we report a carbene-mediated functionalization of the sp³-carbon of the 3-methyl group on 2-formyl-indoles (Scheme 1c). The reaction goes through a carbene-catalyzed addition of 2-formyl-3-methylindoles to different reactive ketones. An NHC-bound *o*-quinodimethane is formed as the key intermediate.

Substituted 3,4-dihydropyrano[3,4-*b*]indoles (indole-derived lactones) are afforded as the products.

RESULTS AND DISCUSSION

We chose the 2,2,2-trifluoroacetophenone **2a** as a model electrophile and tested the model reaction using different NHC catalysts. Imidazolium NHC precursor **A** was not efficient for this transformation (Table 1, entry 1). Triazolium-type NHC precursors could give the product in good isolated yields, and NHC precursor **C**¹⁰ bearing a morpholine motif behaved better (entries 2 and 3). Due to the switching of the conjugate anion in NHC precursors from BF₄[−] to Cl[−], the product yield could be consistently improved (entry 4 vs entry 3). A variety of organic and inorganic bases could be used in this reaction (entries 5–7). THF was found to be the best solvent for this reaction; other organic solvents that we tested gave trace amounts of the desired product (entries 8–10).


With an optimized condition in hand, we next examined the substrate scope. Examples of substituted 2-formyl-3-methylindoles **1** were first tested (Table 2). Both electron-donating groups (**3b** to **3h**) and electron-withdrawing groups (**3i** to **3l**) were well-tolerated. All of the desired hydropyrano[3,4-*b*]indole products were isolated in good to excellent yields. Attempts were also made to synthesize other 3-substituted indoles, but they were not successfully generated.

Meanwhile, different activated ketones **2** were also investigated (Table 3). The benzene ring of the 2,2,2-trifluoroacetophenone **2a** could be changed from substituted benzene groups to heterocyclic aromatic groups, and the corresponding products were isolated in good yields (**3m**–**3q**). To our delight, α -ketoesters also worked very well in this process, and 3,4-dihydropyrano[3,4-*b*]indole-3-carboxylate


Received: September 26, 2017

Published: November 13, 2017




Scheme 1. Hydropyrano[3,4-*b*]indoles and NHC-Catalyzed Functionalization of Enals and Aryl Aldehydesa) Bio-active molecules containing hydropyrano[3,4-*b*]indole:

b) NHC-mediated functionalization of enals and aryl aldehydes:



c) This work:

Table 1. Optimization of Reaction Conditions^a

1a Boc

2a

NHC (30 mol%)
base (30 mol%)
4 (120 mol%)
solvent, RT, 24 h

3a Boc

A

B


C: X = BF₄
D: X = Cl

entry	NHC	base	solvent	yield (%) ^b
1	A	Cs ₂ CO ₃	THF	<5
2	B	Cs ₂ CO ₃	THF	63
3	C	Cs ₂ CO ₃	THF	70
4	D	Cs ₂ CO ₃	THF	95
5	D	K ₂ CO ₃	THF	88
6	D	DMAP	THF	82
7	D	DBU	THF	76
8	D	Cs ₂ CO ₃	CH ₂ Cl ₂	<5
9	D	Cs ₂ CO ₃	EtOAc	10
10	D	Cs ₂ CO ₃	CH ₃ CN	<5


^aReaction conditions: 1a (0.12 mmol), 2a (0.10 mmol), NHC (0.03 mmol), base (0.03 mmol), 4 (0.12 mmol), solvent (2.0 mL), rt, 24 h.

^bIsolated yield of 3a.

compounds were formed smoothly as the products (3r–3t). However, the highly activated ketone 2u bearing both an ester

Table 2. Examples of 2-Formyl-3-methylindoles^a

^aReaction conditions as stated in Table 1, entry 4. Yields are isolated yields after purification by column chromatography.


Table 3. Examples of Substituted Ketones^a

^aReaction conditions as stated in Table 1, entry 4. Yields are isolated yields after purification by column chromatography.

group and a trifluoromethyl group on the carbonyl carbon gave the desired product **3u** in a poor yield.

We also examined chiral NHC catalysts for this reaction. Preliminary studies showed that the use of chiral NHC catalysts **E**¹¹ and **F**¹² could lead to products with low to moderate enantioselectivities (Scheme 2). Additional studies using known


Scheme 2. Enantioselective [4+2] Reactions Using Chiral NHC Catalysts



chiral NHC catalysts did not lead to acceptable enantioselectivities. It appears either new NHC catalysts or the introduction of a cocatalyst is necessary to achieve good stereoselectivities for this reaction.

The plausible catalytic cycle is summarized in Scheme 3. NHC catalyst first attacks the 3-methylindole-2-carboxaldehyde

Scheme 3. Proposed Reaction Mechanism



1a and then goes through proton transfer to afford the Breslow intermediate **I**. Breslow intermediate **I** is then oxidized by **4** to form the acyl azolium intermediate **II**, which is easily deprotonated to afford the *o*-quinodimethane intermediate **III**. Intermediate **III** could react with trifluoroacetophenone **2a** through either a stepwise Michael reaction/cycloaddition sequence or a concerted [4+2] annulation reaction to give intermediate **IV**, which then regenerates the NHC catalyst and yields the final product **3a**.

In summary, we have developed a carbene-catalyzed functionalization of the 3-methyl sp^3 -carbon attached to 2-formyl-indoles. The reaction proceeds through an NHC-bound *o*-quinodimethane as the key intermediate, which is generated

from 2-formyl-3-methylindoles under oxidative conditions. 2,2,2-Trifluoroacetophenones and α -ketoesters are found to be effective substrates to produce substituted 3,4-dihydropyrano[3,4-*b*]indole compounds in good to excellent yields. Additional studies on aromatic compound functionalizations and evaluation on the biological activities of the products are in progress.

EXPERIMENTAL SECTION

General Information. Commercially available materials purchased from J&K or Aladdin were used as received. DCM was dried over a Pure Solv solvent purification system. THF was distilled over sodium. Ethyl acetate was dried over 4 Å molecular sieves prior use. Unless otherwise specified, all reactions were carried out under an atmosphere of nitrogen in a 10 mL Schlenk tube. Proton nuclear magnetic resonance (^1H NMR) spectra were recorded on a Bruker (400 MHz) spectrometer and JEOL-ECX-500 (500 MHz) spectrometer. Chemical shifts were recorded in parts per million (ppm, δ) relative to tetramethylsilane ($\delta = 0.00$) or chloroform ($\delta = 7.26$, singlet). ^1H NMR splitting patterns are designated as singlet (s), doublet (d), triplet (t), quartet (q), dd (doublet of doublets), or m (multiplets). All first-order splitting patterns were assigned on the basis of the appearance of the multiplet. Splitting patterns that could not be easily interpreted are designated as multiplet (m) or broad (br). Carbon nuclear magnetic resonance (^{13}C NMR) spectra were recorded on a Bruker (400 MHz) (101 MHz) spectrometer and JEOL-ECX-500 (126 MHz) spectrometer. The melting points (mp) of the title compounds were determined when left untouched on an XT-4-MP apparatus from Beijing Tech. Instrument Co. (Beijing, China). High-resolution mass spectral analysis (HRMS) was performed on a quadrupole/electrostatic field orbitrap mass spectrometer. X-ray crystallography analysis was performed on Bruker X8 APEX X-ray diffractometer. Analytical thin-layer chromatography (TLC) was carried out on Merck 60 F254 precoated silica gel plates (0.2 mm thickness). Visualization was performed using a UV lamp or potassium permanganate stain. The aldehyde substrates **1** were synthesized through reported procedures.¹³

General Procedure for the Catalytic Reactions of Aldehydes **1 with Trifluoromethyl Aryl Ketones **2**.** To a dry Schlenk reaction tube equipped with a magnetic stir bar were added indolealdehyde **1a** (0.12 mmol), triazolium salt **D** (8.4 mg, 0.03 mmol), oxidant **4** (49 mg, 0.12 mmol), and Cs_2CO_3 (10 mg, 0.03 mmol). The Schlenk tube was then closed with a septum, evacuated, and refilled with N_2 . Freshly distilled anhydrous THF (2 mL) was added, followed by the injection of trifluoromethyl aryl ketone **2a** (0.1 mmol). The mixture was stirred at 25 °C for 24 h. After completion of the reaction, monitored by TLC and ^1H NMR, the solvent was removed under reduced pressure, and the residue was purified via column chromatography on silica gel with hexane/EtOAc (20:1) as an eluent to afford product **3a**.

General Procedure for the Enantioselective Catalytic Reactions of Aldehyde **1a with Trifluoromethyl Aryl Ketones **2a**.** To a dry Schlenk reaction tube equipped with a magnetic stir bar were added indolealdehyde **1a** (0.12 mmol), chiral pre-NHC **E** (12.6 mg, 0.03 mmol) or chiral pre-NHC **F** (11.0 mg, 0.03 mmol), oxidant **4** (49 mg, 0.12 mmol), and Cs_2CO_3 (10 mg, 0.03 mmol). The Schlenk tube was then closed with a septum, evacuated, and refilled with N_2 . Freshly distilled anhydrous THF (2 mL) was added, followed by the injection of trifluoromethyl aryl ketone **2a** (0.1 mmol). The mixture was stirred at rt for 24 h. After completion of the reactions, monitored by TLC and ^1H NMR, the solvent was removed under reduced pressure, and the residue was purified via column chromatography on silica gel with hexane/EtOAc (20:1) as an eluent to afford the products **3**.

tert-Butyl-3-formyl-3-methyl-1H-indole-1-carboxylate (1a**):** white solid; 0.6 g, 22% yield; mp 68–69 °C; ^1H NMR (400 MHz, CDCl_3) δ 10.44 (s, 1H), 8.14 (d, $J = 8.5$ Hz, 1H), 7.65 (d, $J = 7.1$ Hz, 1H), 7.49 (t, $J = 7.2$ Hz, 1H), 7.31 (t, $J = 7.1$ Hz, 1H), 2.57 (s, 3H), 1.69 (s, 9H); ^{13}C NMR (101 MHz, CDCl_3) δ 185.3, 150.0, 136.6, 132.8, 129.4, 128.4, 128.3, 123.4, 121.0, 115.9, 85.1, 28.2, 10.1 ppm; HRMS (ESI-

TOF) m/z $[M + Na]^+$ calcd for $C_{15}H_{17}NO_3Na^+$ 282.1100, found 282.1100.

tert-Butyl-2-formyl-3,6-dimethyl-1H-indole-1-carboxylate (1b): white solid; 0.52 g, 19% yield; mp 90–91 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.39 (s, 1H), 7.99 (s, 1H), 7.53 (d, J = 8.1 Hz, 1H), 7.14 (d, J = 7.3 Hz, 1H), 2.55 (s, 3H), 2.51 (s, 3H), 1.69 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.1, 150.0, 139.1, 137.2, 132.5, 128.8, 127.2, 125.1, 120.7, 116.0, 85.0, 28.2, 22.4, 10.2 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{16}H_{19}NO_3Na^+$ 296.1257, found 296.1257.

tert-Butyl-2-formyl-3,5-dimethyl-1H-indole-1-carboxylate (1c): white solid; 0.43 g, 16% yield; mp 88–89 °C; 1H NMR (500 MHz, $CDCl_3$) δ 10.39 (s, 1H), 7.97 (d, J = 8.6 Hz, 1H), 7.37 (s, 1H), 7.27 (dd, J = 8.6, 1.4 Hz, 1H), 2.50 (s, 3H), 2.44 (s, 3H), 1.68 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 185.4, 150.0, 134.9, 133.0, 130.1, 129.5, 128.1, 120.7, 115.6, 84.9, 28.2, 21.3, 10.1 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{16}H_{19}NO_3Na^+$ 296.1257, found 296.1255.

tert-Butyl-2-formyl-3,4-dimethyl-1H-indole-1-carboxylate (1d): red solid; 0.54 g, 20% yield; mp 89–90 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.32 (s, 1H), 8.01 (d, J = 8.5 Hz, 1H), 7.38–7.28 (m, 1H), 7.01 (d, J = 7.3 Hz, 1H), 2.73 (s, 3H), 2.71 (s, 3H), 1.67 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 184.9, 149.8, 137.3, 134.0, 132.7, 128.9, 128.0, 127.8, 125.4, 113.5, 85.1, 28.1, 20.8, 12.6 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{16}H_{19}NO_3Na^+$ 296.1257, found 296.1255.

tert-Butyl-6-(tert-butyl)-2-formyl-3-methyl-1H-indole-1-carboxylate (1e): yellow solid; 0.81 g, 21% yield; mp 84–85 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.44 (s, 1H), 8.19 (s, 1H), 7.58 (d, J = 8.4 Hz, 1H), 7.40 (dd, J = 8.4, 1.7 Hz, 1H), 2.56 (s, 3H), 1.71 (s, 9H), 1.41 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.3, 152.4, 150.2, 136.9, 133.0, 128.5, 127.2, 121.6, 120.5, 112.4, 84.7, 35.5, 31.6, 28.2, 10.2 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{19}H_{25}NO_3Na^+$ 338.1726, found 338.1722.

tert-Butyl-2-formyl-5-methoxy-3-methyl-1H-indole-1-carboxylate (1f): white solid; 0.51 g, 13% yield; mp 117–118 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.43 (s, 1H), 8.02 (d, J = 9.1 Hz, 1H), 7.11 (dd, J = 9.2, 2.6 Hz, 1H), 7.01 (d, J = 2.5 Hz, 1H), 3.88 (s, 3H), 2.54 (s, 3H), 1.69 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.4, 156.2, 149.9, 133.4, 131.3, 130.1, 127.9, 118.2, 116.9, 102.0, 85.0, 55.6, 28.2, 10.2 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{16}H_{19}NO_4Na^+$ 312.1206, found 312.1203.

tert-Butyl-2-formyl-6-methoxy-3-methyl-1H-indole-1-carboxylate (1g): white solid; 0.45 g, 11% yield; mp 88–89 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.37 (s, 1H), 7.68 (s, 1H), 7.52 (d, J = 8.7 Hz, 1H), 6.94 (d, J = 7.7 Hz, 1H), 3.90 (s, 3H), 2.55 (s, 3H), 1.69 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 184.6, 161.0, 150.1, 138.3, 132.2, 129.5, 123.2, 121.8, 113.7, 98.9, 84.9, 55.6, 28.2, 10.3 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{16}H_{19}NO_4Na^+$ 312.1206, found 312.1200.

tert-Butyl-2-formyl-3-methyl-6-phenyl-1H-indole-1-carboxylate (1h): white solid; 0.45 g, 23% yield; mp 144–145 °C; 1H NMR (500 MHz, $CDCl_3$) δ 10.45 (s, 1H), 8.42 (s, 1H), 7.69 (dd, J = 15.1, 7.6 Hz, 3H), 7.58 (dd, J = 8.2, 1.5 Hz, 1H), 7.48 (t, J = 7.6 Hz, 2H), 7.38 (t, J = 7.4 Hz, 1H), 2.59 (s, 3H), 1.70 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 185.3, 150.1, 141.9, 141.3, 137.3, 133.3, 129.0, 128.7, 128.5, 127.6, 127.5, 123.1, 121.4, 114.6, 85.3, 28.3, 10.3 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{21}H_{21}NO_3Na^+$ 358.1413, found 358.1409.

tert-Butyl-5-fluoro-2-formyl-3-methyl-1H-indole-1-carboxylate (1i): white solid; 0.63 g, 16% yield; mp 117–118 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.42 (s, 1H), 8.10 (dd, J = 9.2, 4.5 Hz, 1H), 7.39–7.12 (m, 2H), 2.51 (s, 3H), 1.69 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.2, 160.5, 158.1, 149.7, 133.9, 132.8, 130.3 (d, J_{C-F} = 9.2 Hz), 127.4 (d, J_{C-F} = 4.5 Hz), 117.2 (d, J_{C-F} = 8.7 Hz), 116.6 (d, J_{C-F} = 25.4 Hz), 106.0 (d, J_{C-F} = 23.3 Hz), 85.5, 28.2, 10.1 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{15}H_{16}NO_3FNa^+$ 300.1006, found 300.1004.

tert-Butyl-6-fluoro-2-formyl-3-methyl-1H-indole-1-carboxylate (1j): red solid; 0.72 g, 18% yield; mp 95–96 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.41 (s, 1H), 7.84 (dd, J = 10.5, 2.3 Hz, 1H), 7.58 (dd, J = 8.7, 5.5 Hz, 1H), 7.15–6.95 (m, 1H), 2.54 (s, 3H), 1.70 (s, 9H); ^{13}C

NMR (101 MHz, $CDCl_3$) δ 184.8, 164.5, 162.1, 149.7, 137.0 (d, J_{C-F} = 13.2 Hz), 133.3 (d, J_{C-F} = 4.0 Hz), 128.2, 125.7, 122.2 (d, J_{C-F} = 10.4 Hz), 112.2 (d, J_{C-F} = 24.9 Hz), 103.1 (d, J_{C-F} = 28.9 Hz), 85.6, 28.1, 10.2 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{15}H_{16}NO_3FNa^+$ 300.1006, found 300.1002.

tert-Butyl-5-bromo-2-formyl-3-methyl-1H-indole-1-carboxylate (1k): white solid; 0.48 g, 19% yield; mp 125–126 °C; 1H NMR (500 MHz, $CDCl_3$) δ 10.41 (s, 1H), 8.01 (d, J = 8.8 Hz, 1H), 7.77 (d, J = 1.7 Hz, 1H), 7.56 (dd, J = 9.0, 1.8 Hz, 1H), 2.51 (s, 3H), 1.68 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.4, 150.0, 136.6, 132.9, 129.4, 128.5, 128.4, 123.4, 121.1, 116.0, 85.2, 28.2, 10.1 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{15}H_{16}NO_3BrNa^+$ 360.0205, found 360.0201.

tert-Butyl-6-chloro-2-formyl-3-methyl-1H-indole-1-carboxylate (1l): white solid; 0.93 g, 16% yield; mp 111–112 °C; 1H NMR (400 MHz, $CDCl_3$) δ 10.40 (s, 1H), 8.17 (d, J = 1.5 Hz, 1H), 7.54 (d, J = 8.5 Hz, 1H), 7.27 (dd, J = 8.5, 1.8 Hz, 1H), 2.53 (s, 3H), 1.70 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 185.0, 149.5, 136.8, 134.4, 133.1, 127.8, 127.8, 124.1, 121.8, 116.2, 85.8, 28.1, 10.0 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{15}H_{16}NO_3ClNa^+$ 316.0710, found 316.0711.

tert-Butyl-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3a): yellow solid; mp 128–129 °C; 41 mg, 95% yield; 1H NMR (500 MHz, $CDCl_3$) δ 8.07 (d, J = 8.4 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.58 (dd, J = 6.4, 2.8 Hz, 2H), 7.47 (t, J = 7.5 Hz, 1H), 7.37–7.32 (m, 3H), 7.30 (t, J = 7.4 Hz, 1H), 3.88 (d, J = 17.3 Hz, 1H), 3.74 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 154.1, 148.6, 139.9, 133.8, 129.7, 129.4, 128.7, 127.6, 126.6, 125.0, 124.4, 123.8, 122.2, 120.6, 115.3, 84.9, 83.8 (q, J = 30.6 Hz), 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{23}H_{20}NO_4F_3Na^+$ 454.1236, found 454.1231.

tert-Butyl-7-methyl-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3b): yellow solid; mp 177–178 °C; 39 mg, 89% yield; 1H NMR (500 MHz, $CDCl_3$) δ 7.91 (s, 1H), 7.60–7.54 (m, 2H), 7.48 (d, J = 8.1 Hz, 1H), 7.35–7.30 (m, 3H), 7.14 (d, J = 8.1 Hz, 1H), 3.84 (d, J = 17.3 Hz, 1H), 3.72 (d, J = 17.3 Hz, 1H), 2.47 (s, 3H), 1.58 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 154.2, 148.8, 140.4, 133.9, 129.6, 128.7, 127.8, 126.6, 125.5, 124.4, 123.2, 122.8, 122.2, 120.2, 115.3, 84.8, 83.7 (q, J = 30.9 Hz), 27.6, 26.2, 22.4 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{24}H_{22}NO_4F_3Na^+$ 468.1393, found 468.1391.

tert-Butyl-6-methyl-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3c): yellow solid; mp 145–146 °C; 36 mg, 81% yield; 1H NMR (500 MHz, $CDCl_3$) δ 7.95 (d, J = 8.6 Hz, 1H), 7.57 (dd, J = 6.1, 3.1 Hz, 2H), 7.39 (s, 1H), 7.35–7.32 (m, 3H), 7.31 (dd, J = 8.7, 1.4 Hz, 1H), 3.85 (d, J = 17.3 Hz, 1H), 3.71 (d, J = 17.2 Hz, 1H), 2.45 (s, 3H), 1.58 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 154.1, 148.7, 138.2, 133.9, 133.5, 131.1, 129.7, 128.7, 127.4, 126.6, 125.2, 124.4, 123.7, 120.1, 115.0, 84.7, 83.7 (q, J = 30.2 Hz), 27.6, 26.2, 21.3 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{24}H_{22}NO_4F_3Na^+$ 468.1393, found 468.1395.

tert-Butyl-5-methyl-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3d): white solid; mp 153–154 °C; 39 mg, 89% yield; 1H NMR (400 MHz, $CDCl_3$) δ 7.91 (d, J = 8.5 Hz, 1H), 7.58 (dd, J = 6.5, 2.7 Hz, 2H), 7.38–7.30 (m, 4H), 7.03 (d, J = 7.3 Hz, 1H), 4.14 (d, J = 17.4 Hz, 1H), 3.90 (d, J = 17.4 Hz, 1H), 2.71 (s, 3H), 1.57 (s, 9H); ^{13}C NMR (126 MHz, $CDCl_3$) δ 154.2, 148.7, 140.2, 133.8, 132.7, 129.7, 129.1, 128.8, 127.5, 126.7, 125.0, 124.1, 123.6, 122.2, 112.8, 84.8, 83.5 (q, J = 30.2 Hz), 28.4, 27.6, 20.0 ppm; HRMS (ESI-TOF) m/z $[M + Na]^+$ calcd for $C_{24}H_{22}NO_4F_3Na^+$ 468.1393, found 468.1391.

tert-Butyl-7-(tert-butyl)-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3e): white solid; mp 116–117 °C; 47 mg, 97% yield; 1H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1H), 7.57 (dd, J = 6.0, 3.0 Hz, 2H), 7.54 (d, J = 8.5 Hz, 1H), 7.40 (dd, J = 8.5, 1.5 Hz, 1H), 7.37–7.30 (m, 3H), 3.86 (d, J = 17.3 Hz, 1H), 3.72 (d, J = 17.3 Hz, 1H), 1.60 (s, 9H), 1.36 (s, 9H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 154.0, 153.6, 148.7, 140.2, 133.9, 129.5, 128.6, 127.5, 126.5, 124.6, 123.5, 122.7, 122.1, 119.9, 111.7, 84.6, 83.6 (q, J =

61.0, 30.4 Hz), 35.6, 31.4, 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₇H₂₈NO₄F₃Na⁺ 510.1862, found 510.1861.

tert-Butyl-6-methoxy-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3f): white solid; mp 159–160 °C; 38 mg, 83% yield; ¹H NMR (500 MHz, CDCl₃) δ 7.97 (d, J = 9.2 Hz, 1H), 7.58 (dd, J = 6.1, 2.7 Hz, 2H), 7.38–7.31 (m, 3H), 7.10 (dd, J = 9.2, 2.5 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 3.87 (s, 3H), 3.84 (d, J = 17.3 Hz, 1H), 3.72 (d, J = 17.3 Hz, 1H), 1.58 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 156.6, 154.0, 148.7, 134.8, 133.9, 129.7, 128.7, 127.1, 126.6, 125.6, 124.0, 122.2, 119.4, 116.4, 101.5, 84.8, 83.7 (q, 33.1 Hz), 55.8, 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₄H₂₂NO₅F₃Na⁺ 484.1342, found 484.1349.

tert-Butyl-7-methoxy-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3g): yellow solid; mp 177–178 °C; 41 mg, 89% yield; ¹H NMR (400 MHz, CDCl₃) δ 7.58 (dd, J = 8.6, 2.8 Hz, 3H), 7.47 (d, J = 8.8 Hz, 1H), 7.35–7.31 (m, 3H), 6.93 (dd, J = 8.8, 2.3 Hz, 1H), 3.86 (s, 3H), 3.82 (d, J = 17.3 Hz, 1H), 3.70 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 161.8, 154.0, 149.0, 141.7, 134.0, 129.6, 128.7, 128.3, 126.5, 124.4, 122.4, 121.3, 118.7, 114.7, 97.8, 84.8, 83.5 (q, J = 30.4 Hz), 55.8, 27.6, 26.3 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₄H₂₂NO₅F₃Na⁺ 484.1342, found 484.1349.

tert-Butyl-1-oxo-3,7-diphenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3h): white solid; mp 86–87 °C; 48 mg, 96% yield; ¹H NMR (400 MHz, CDCl₃) δ 8.33 (s, 1H), 7.65 (dd, J = 16.6, 7.8 Hz, 3H), 7.61–7.54 (m, 3H), 7.45 (t, J = 7.5 Hz, 2H), 7.40–7.32 (m, 4H), 3.90 (d, J = 17.3 Hz, 1H), 3.77 (d, J = 17.3 Hz, 1H), 1.60 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 153.9, 148.6, 142.9, 140.7, 140.5, 133.8, 129.6, 128.9, 128.7, 127.8, 127.5, 127.4, 126.5, 124.6, 124.0, 123.5, 121.8, 120.7, 113.7, 85.0, 83.8 (q, J = 30.7 Hz), 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₉H₂₄NO₄F₃Na⁺ 530.1549, found 530.1551.

tert-Butyl-6-fluoro-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3i): yellow solid; mp 161–162 °C; 39 mg, 86% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.06 (dd, J = 9.2, 4.4 Hz, 1H), 7.57 (s, 2H), 7.38–7.33 (m, 3H), 7.28–7.23 (m, 1H), 7.21 (dd, J = 9.1, 2.4 Hz, 1H), 3.82 (d, J = 17.3 Hz, 1H), 3.73 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 160.5, 158.5, 153.8, 148.4, 136.2, 133.7, 129.8, 128.8, 126.8 (d, J_{C-F} = 4.2 Hz), 126.5, 125.6 (d, J_{C-F} = 9.7 Hz), 125.0, 124.3, 122.1, 117.7 (d, J_{C-F} = 25.6 Hz), 116.8 (d, J_{C-F} = 8.9 Hz), 105.6 (d, J_{C-F} = 24.0 Hz), 85.3, 83.8 (q, J = 30.5 Hz), 27.6, 26.1 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₄Na⁺ 472.1142, found 472.1140.

tert-Butyl-7-fluoro-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3j): white solid; mp 159–160 °C; 38 mg, 85% yield; ¹H NMR (500 MHz, CDCl₃) δ 7.80 (dd, J = 10.0, 2.3 Hz, 1H), 7.56 (dd, J = 8.7, 5.3 Hz, 3H), 7.37–7.31 (m, 3H), 7.07 (td, J = 8.8, 2.3 Hz, 1H), 3.84 (d, J = 17.3 Hz, 1H), 3.73 (d, J = 17.4 Hz, 1H), 1.58 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 164.8, 162.8, 153.7, 148.4, 140.5 (d, J_{C-F} = 13.3 Hz), 133.7, 129.8, 128.8, 127.5, 126.5, 124.2 (d, J_{C-F} = 34.3 Hz), 121.8 (d, J_{C-F} = 10.5 Hz), 121.4, 113.0 (d, J_{C-F} = 25.2 Hz), 102.6 (d, J_{C-F} = 28.7 Hz), 85.4, 83.7 (q, J = 30.2 Hz), 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₄Na⁺ 472.1142, found 472.1140.

tert-Butyl-6-bromo-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3k): white solid; mp 132–134 °C; 38 mg, 75% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.08 (d, J = 8.3 Hz, 1H), 7.61 (d, J = 6.1, 1H), 7.58 (dd, J = 6.1, 3.1 Hz, 2H), 7.52–7.45 (m, 1H), 7.33 (t, 3H), 3.88 (d, J = 17.3 Hz, 1H), 3.75 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 154.0, 148.6, 139.9, 133.8, 129.7, 129.4, 128.7, 127.6, 126.6, 125.0, 124.4, 123.8, 122.1, 120.6, 115.4, 85.0, 83.8 (q, J = 30.4 Hz), 27.6, 26.2 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₃BrNa⁺ 532.0341, found 532.0340.

tert-Butyl-7-chloro-1-oxo-3-phenyl-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3l): white solid; mp 158–160 °C; 41 mg, 88% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.13 (d, J = 1.6 Hz, 1H), 7.58–7.54 (m, 2H), 7.53 (d, J = 8.5 Hz, 1H), 7.37–7.32 (m, 3H), 7.29 (dd, J = 8.5, 1.7 Hz, 1H), 3.85 (d, J = 17.3 Hz, 1H), 3.74 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ¹³C NMR (126

MHz, CDCl₃) δ 153.8, 148.3, 140.1, 135.6, 133.6, 129.8, 128.8, 127.2, 126.5, 124.7, 124.2, 123.5, 122.1, 121.3, 115.6, 85.5, 83.8 (q, J = 30.7 Hz), 27.6, 26.1 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₃ClNa⁺ 488.0846, found 488.0844.

tert-Butyl-1-oxo-3-(p-tolyl)-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3m): white solid; mp 158–159 °C; 39 mg, 89% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.08 (d, J = 8.5 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.54–7.40 (m, 3H), 7.31 (t, J = 7.5 Hz, 1H), 7.13 (d, J = 8.1 Hz, 2H), 3.85 (d, J = 17.2 Hz, 1H), 3.72 (d, J = 17.2 Hz, 1H), 2.29 (s, 3H), 1.60 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 154.1, 148.7, 139.8, 139.7, 130.8, 129.4, 129.3, 127.6, 126.4, 125.0, 124.4, 123.8, 122.2, 120.6, 115.4, 84.9, 83.8 (q, J = 30.3 Hz), 27.7, 26.2, 21.1 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₄H₂₂NO₄F₃Na⁺ 468.1393, found 468.1390.

tert-Butyl-3-(4-fluorophenyl)-1-oxo-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3n): yellow solid; mp 147–149 °C; 33 mg, 73% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.09 (d, J = 8.5 Hz, 1H), 7.62 (d, J = 8.0 Hz, 1H), 7.56 (dd, J = 8.7, 5.0 Hz, 2H), 7.51 (t, J = 7.8 Hz, 1H), 7.33 (t, J = 7.5 Hz, 1H), 7.03 (t, J = 8.6 Hz, 2H), 3.84 (d, J = 17.3 Hz, 1H), 3.76 (d, J = 17.3 Hz, 1H), 1.59 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 164.6, 162.1, 153.7, 148.5, 139.8, 129.6 (d, J_{C-F} = 3.4 Hz), 129.5, 128.5 (d, J_{C-F} = 8.5 Hz), 127.4, 124.8, 123.8, 123.5, 120.4, 115.9, 115.7, 115.3, 85.0, 83.4 (q, J = 30.9 Hz), 27.6, 26.1 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₄Na⁺ 472.1142, found 472.1140.

tert-Butyl-3-(4-chlorophenyl)-1-oxo-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3o): white solid; mp 170–171 °C; 40 mg, 87% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.09 (d, J = 8.6 Hz, 1H), 7.61 (d, J = 7.9 Hz, 1H), 7.51 (t, J = 9.1 Hz, 3H), 7.32 (t, J = 8.5 Hz, 3H), 3.83 (d, J = 17.3 Hz, 1H), 3.75 (d, J = 17.3 Hz, 1H), 1.60 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 153.7, 148.5, 139.9, 136.0, 132.5, 129.6, 129.0, 128.0, 127.3, 124.8, 123.9, 123.5, 121.9, 120.5, 115.4, 85.1, 83.5 (d, J = 31.0 Hz), 27.7, 26.1 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₃ClNa⁺ 488.0846, found 488.0841.

tert-Butyl-3-(4-bromophenyl)-1-oxo-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3p): white solid; mp 169–170 °C; 41 mg, 89% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.09 (d, J = 8.5 Hz, 1H), 7.61 (d, J = 7.7 Hz, 1H), 7.55–7.39 (m, 5H), 7.32 (t, J = 7.5 Hz, 1H), 3.82 (d, J = 17.3 Hz, 1H), 3.75 (d, J = 17.3 Hz, 1H), 1.60 (s, 9H); ¹³C NMR (101 MHz, CDCl₃) δ 153.6, 148.4, 139.8, 133.0, 131.9, 129.5, 128.2, 127.2, 124.8, 124.2, 123.8, 123.4, 121.5, 120.4, 115.4, 85.1, 83.4 (q, J = 30.7 Hz), 27.6, 26.0 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₃H₁₉NO₄F₃BrNa⁺ 532.0341, found 532.0348.

tert-Butyl-1-oxo-3-(thiophen-2-yl)-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-b]indole-9(1H)-carboxylate (3q): white solid; mp 149–150 °C; 39 mg, 89% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.12 (d, J = 8.4 Hz, 1H), 7.63 (d, J = 7.9 Hz, 1H), 7.51 (t, J = 7.9 Hz, 1H), 7.34 (t, J = 7.6 Hz, 1H), 7.30 (d, J = 5.1 Hz, 1H), 7.23 (d, J = 3.6 Hz, 1H), 6.95 (t, J = 4.3 Hz, 1H), 3.76 (s, 2H), 1.60 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 153.4, 148.7, 139.9, 137.5, 129.5, 128.0, 127.4, 127.3, 125.0, 123.9, 123.2, 121.6, 120.5, 115.4, 85.1, 82.5 (q, J = 31.9 Hz), 27.8, 27.7 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₁H₁₈NO₄F₃SNa⁺ 460.0800, found 460.0800.

9-tert-Butyl-3-ethyl-3-(4-chlorophenyl)-1-oxo-3,4-dihydropyrano[3,4-b]indole-3,9(1H)-dicarboxylate (3r): white solid; mp 121–122 °C; 41 mg, 87% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.19 (d, J = 8.5 Hz, 1H), 7.71 (d, J = 8.6 Hz, 2H), 7.65 (d, J = 7.9 Hz, 1H), 7.56–7.50 (m, 1H), 7.42 (d, J = 8.6 Hz, 2H), 7.34 (t, J = 7.5 Hz, 1H), 4.19 (d, J = 17.0 Hz, 1H), 4.12 (q, J = 7.1 Hz, 2H), 3.33 (d, J = 17.0 Hz, 1H), 1.67 (s, 9H), 1.12 (t, J = 7.1 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 170.3, 155.4, 148.9, 139.9, 135.8, 135.1, 129.6, 129.4, 129.0, 126.7, 124.9, 123.9, 123.8, 121.0, 115.5, 85.0, 84.6, 62.9, 31.4, 27.8, 14.0 ppm; HRMS (ESI-TOF) m/z [M + Na]⁺ calcd for C₂₅H₂₄NO₆ClNa⁺ 492.1184, found 492.1189.

9-tert-Butyl-3-methyl-1-oxo-3-phenyl-3,4-dihydropyrano[3,4-b]indole-3,9(1H)-dicarboxylate (3s): white solid; mp 158–160 °C; 35 mg, 83% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.18 (d, J = 8.5 Hz, 1H), 7.77 (d, J = 7.3 Hz, 2H), 7.66 (d, J = 7.8 Hz, 1H), 7.54 (t, J = 7.8

H₂, 1H), 7.49–7.38 (m, 3H), 7.34 (t, *J* = 7.5 Hz, 1H), 4.21 (d, *J* = 17.2 Hz, 1H), 3.67 (s, 3H), 3.40 (d, *J* = 15.7 Hz, 1H), 1.67 (s, 9H); ¹³C NMR (126 MHz, CDCl₃) δ 171.2, 155.7, 149.0, 140.1, 137.0, 130.0, 129.3, 129.1, 128.8, 125.2, 125.0, 124.2, 123.7, 121.1, 115.4, 85.2, 84.9, 53.5, 31.5, 27.8 ppm; HRMS (ESI-TOF) *m/z* [M + Na]⁺ calcd for C₂₄H₂₃NO₆Na⁺ 444.1417, found 444.1419.

9-tert-Butyl-3-ethyl-1-oxo-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-*b*]indole-3,9(1*H*)-dicarboxylate (3t): yellow solid; mp 157–158 °C; 38 mg, 88% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.18 (d, *J* = 8.5 Hz, 1H), 7.79–7.73 (m, 2H), 7.66 (d, *J* = 7.9 Hz, 1H), 7.57–7.48 (m, 1H), 7.46–7.41 (m, 2H), 7.41–7.36 (m, 1H), 7.34 (t, *J* = 7.6 Hz, 1H), 4.21 (d, *J* = 17.1 Hz, 1H), 4.11 (q, *J* = 7.1 Hz, 2H), 3.37 (d, *J* = 17.1 Hz, 1H), 1.67 (s, 9H), 1.12 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 170.6, 155.7, 149.0, 139.9, 137.2, 129.9, 129.3, 129.0, 128.8, 125.2, 125.0, 124.1, 123.7, 121.0, 115.4, 85.1, 84.9, 62.7, 31.4, 27.8, 14.0 ppm; HRMS (ESI-TOF) *m/z* [M + Na]⁺ calcd for C₂₅H₂₅NO₆Na⁺ 458.1574, found 458.1579.

9-tert-Butyl-3-ethyl-1-oxo-3-(trifluoromethyl)-3,4-dihydropyrano[3,4-*b*]indole-3,9(1*H*)-dicarboxylate (3u): white solid; mp 109–110 °C; 17 mg, 40% yield; ¹H NMR (500 MHz, CDCl₃) δ 8.19 (d, *J* = 8.5 Hz, 1H), 7.62 (d, *J* = 8.0 Hz, 1H), 7.56 (dd, *J* = 11.9, 4.4 Hz, 1H), 7.36 (t, *J* = 7.4 Hz, 1H), 4.25 (q, *J* = 7.1 Hz, 2H), 3.89 (d, *J* = 17.1 Hz, 1H), 3.45 (d, *J* = 17.0 Hz, 1H), 1.66 (s, 9H), 1.22 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ 165.2, 152.9, 148.6, 139.9, 129.8, 126.8, 124.5, 124.0, 122.7, 120.9, 115.5, 85.4, 82.8, 82.6, 64.0, 27.7, 25.1, 13.9 ppm; HRMS (ESI-TOF) *m/z* [M + Na]⁺ calcd for C₂₀H₂₀NO₆F₃Na⁺ 450.1134, found 450.1139.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.7b02436.

Spectral data for all new compounds and X-ray crystallography data (PDF)
Crystal data of 3a (CIF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: jhxx.msm@gmail.com.

*E-mail: robinchi@ntu.edu.sg.

ORCID

Song Yang: 0000-0003-1301-3030

Zhichao Jin: 0000-0003-3003-6437

Yonggui Robin Chi: 0000-0003-0573-257X

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We acknowledge financial support by the National Natural Science Foundation of China (no. 21472028), National Key Technologies R&D Program (no. 2014BAD23B01), “Thousand Talent Plan”, The 10 Talent Plan (Shicengci) of Guizhou Province, Guizhou Province Returned Oversea Student Science and Technology Activity Program, Guizhou University (China) and Singapore National Research Foundation (NRF-NRFI2016-06), the Ministry of Education of Singapore (MOE2013-T2-2-003, MOE2016-T2-1-032, and RG108/16), A*STAR Individual Research Grant (A1783c0008), and Nanyang Technological University.

■ REFERENCES

- (1) (a) Lounasmaa, M.; Tolvanen, A. *Nat. Prod. Rep.* **2000**, *17*, 175. (b) Chen, H.; Bai, J.; Fang, Z.-F.; Yu, S.-S.; Ma, S.-G.; Xu, S.; Li, Y.;

Qu, J.; Ren, J.-H.; Li, L.; Si, Y.-K.; Chen, X.-G. *J. Nat. Prod.* **2011**, *74*, 2438 and references cited therein.

- (2) (a) Demerson, C. A.; Abraham, N. A.; Schilling, G.; Martel, R. R.; Pace-Asciak, C.; Humber, L. G. *J. Med. Chem.* **1983**, *26*, 1778. (b) Humber, L. *Med. Res. Rev.* **1987**, *7*, 1. (c) Rabindran, S. K.; He, H.; Singh, M.; Collins, E.; Annable, K. I.; Greeberger, L. M. *Cancer Res.* **1998**, *58*, 5850. (d) Larghi, E. L.; Kaufman, T. S. *Synthesis* **2006**, 187. (e) LaPorte, M. G.; Draper, T. L.; Miller, L. E.; Blackledge, C. W.; Leister, L. K.; Amparo, E.; Hussey, A. R.; Young, D. C.; Chunduru, S. K.; Benetatos, C. A.; Rhodes, G.; Gopalsamy, A.; Herbertz, T.; Burns, C. J.; Condon, S. M. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 2968.

- (3) (a) Demerson, C. A.; Humber, L. G.; Philipp, A. H.; Martel, R. R. *J. Med. Chem.* **1976**, *19*, 391. (b) Katz, A.; Demerson, C. A.; Shaw, C. C.; Asselin, A. A.; Humber, L. G.; Conway, K.; Gavin, G.; Jensen, N. P.; Noureldin, R.; Schmid, J.; Shah, U.; Van Engen, D.; Chau, T.; Weichman, B. J. *Med. Chem.* **1988**, *31*, 1244. (c) Gopalsamy, A.; Lim, K.; Ciszewski, G.; Park, K.; Ellingboe, J. W.; Bloom, J.; Insaf, S.; Upeslakis, J.; Mansour, T. S.; Krishnamurthy, G.; Damarla, M.; Pyatski, Y.; Ho, D.; Howe, A. Y. M.; Orlowski, M.; Feld, B.; O’Connell, J. J. *Med. Chem.* **2004**, *47*, 6603. (d) Demerson, C. A.; Santroch, G.; Humber, L. G.; Charest, M. P. *J. Med. Chem.* **1975**, *18*, 577. (e) Queiroz, M.-J. R. P.; Calhelha, R. C.; Vale-Silva, L. A.; Pinto, E.; São-Jose, N. M. *Eur. J. Med. Chem.* **2009**, *44*, 1893.

- (4) For selected reviews, see: (a) Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *Acc. Chem. Res.* **2008**, *41*, 1013. (b) Dong, Z.; Ren, Z.; Thompson, S. J.; Xu, Y.; Dong, G. *Chem. Rev.* **2017**, *117*, 9333 and references cited therein.

- (5) For recent reviews, see: (a) Bartoli, G.; Bencivenni, G.; Dalpozzo, R. *Chem. Soc. Rev.* **2010**, *39*, 4449. (b) Volla, C. M. R.; Atodiresi, I.; Rueping, M. *Chem. Rev.* **2014**, *114*, 2390. (c) Dalpozzo, R. *Chem. Soc. Rev.* **2015**, *44*, 742. (d) Glinsky-Olivier, N.; Guinchard, X. *Synthesis* **2017**, *49*, 2605. (e) Caruana, L.; Fochi, M.; Bernardi, L. *Synlett* **2017**, *28*, 1530. For selected examples, see (f) Liu, Y.; Nappi, M.; Arceo, E.; Vera, S.; Melchiorre, P. *J. Am. Chem. Soc.* **2011**, *133*, 15212. (g) Jia, Z.-J.; Zhou, Q.; Zhou, Q.-Q.; Chen, P.-Q.; Chen, Y.-C. *Angew. Chem., Int. Ed.* **2011**, *50*, 8638. (h) Liu, Y.; Nappi, M.; Escudero-Adan, E. C.; Melchiorre, P. *Org. Lett.* **2012**, *14*, 1310.

- (6) For selected reviews on NHC catalysis, see: (a) Enders, D.; Niemeier, O.; Henseler, A. *Chem. Rev.* **2007**, *107*, 5606. (b) Flanagan, D. M.; Romanov-Michailidis, F.; White, N. A.; Rovis, T. *Chem. Rev.* **2015**, *115*, 9307 and references cited therein.

- (7) For pioneering works on enolate activation in NHC catalysis, see: (a) Chow, K. Y.-K.; Bode, J. W. *J. Am. Chem. Soc.* **2004**, *126*, 8126. (b) Reynolds, N. T.; Read de Alaniz, J.; Rovis, T. *J. Am. Chem. Soc.* **2004**, *126*, 9518. For pioneering works on homoenolate activation in NHC catalysis, see (c) Burstein, C.; Glorius, F. *Angew. Chem., Int. Ed.* **2004**, *43*, 6205. (d) Sohn, S. S.; Rosen, E. L.; Bode, J. W. *J. Am. Chem. Soc.* **2004**, *126*, 14370. For a pioneering work on oxidative NHC catalysis of enals, see (e) De Sarkar, S.; Studer, A. *Angew. Chem., Int. Ed.* **2010**, *49*, 9266. For pioneering examples on γ - and δ -carbon activation of enals through oxidative NHC catalysis, see: (f) Mo, J.; Chen, X.; Chi, Y. R. *J. Am. Chem. Soc.* **2012**, *134*, 8810. (g) Zhu, T.; Mou, C.; Li, B.; Smetankova, M.; Song, B.-A.; Chi, Y. R. *J. Am. Chem. Soc.* **2015**, *137*, 5658.

- (8) Chen, X.; Yang, S.; Song, B.-A.; Chi, Y. R. *Angew. Chem., Int. Ed.* **2013**, *52*, 11134.

- (9) (a) Janssen-Muller, D.; Singha, S.; Olyschläger, T.; Daniliuc, C. G.; Glorius, F. *Org. Lett.* **2016**, *18*, 4444. (b) Chen, D.-F.; Rovis, T. *Synthesis* **2016**, *49*, 293.

- (10) For a pioneering report on NHC catalysts C and D, see: Chiang, P.-C.; Rommel, M.; Bode, J. W. *J. Am. Chem. Soc.* **2009**, *131*, 8714.

- (11) For a pioneering report on NHC catalyst E, see: Wadamoto, M.; Phillips, E. M.; Reynolds, T. E.; Scheidt, K. A. *J. Am. Chem. Soc.* **2007**, *129*, 10098.

- (12) For a pioneering report on NHC catalyst F, see: He, M.; Struble, J. R.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 8418.

- (13) Tan, W.; Li, X.; Gong, Y.-X.; Ge, M.-D.; Shi, F. *Chem. Commun.* **2014**, *50*, 15901.